

2.1: Functions and Domain and Range

1. If $h \neq 0$ and $f(x) = x^2 - 1$, then $\frac{f(x+h)-f(x)}{h} =$

A) $2x + h$

B) $2x + h + 1$

C) $2x - h - 1$

D) $2x - h$

E) $h - 2$

2. The domain D and the range R of the function $f(x) = 2 - \sqrt{6 - 3x}$ are respectively given by

A) $D = (-\infty, 2]$ and $R = (-\infty, 2]$

B) $D = (-\infty, 2]$ and $R = [2, \infty)$

C) $D = (-\infty, 2]$ and $R = [2, 6]$

D) $D = [2, \infty)$ and $R = [2, \infty)$

E) $D = [2, \infty)$ and $R = (-\infty, 2]$

3. If $f(x) = x^3 - 1$ and $h \neq 0$, then $\frac{f(2+h)-f(2)}{h} =$

A) $h^2 + 6h + 12$

B) $h^2 + 6h + 14$

C) h^2

D) $h^2 - \frac{2}{h}$

E) $h^2 + 6h$

4. The domain of $y = \frac{1}{\sqrt{x-3}}$ in interval notation is:

A) $[0,9) \cup (9, \infty)$

B) $(-\infty, 9) \cup (9, \infty)$

C) $[0, \infty)$

D) $(3, \infty)$

E) $(9, \infty)$

5. The domain D and the range R of the function $f(x) = \frac{\sqrt{4-9x^2}}{2}$ is given by

A) $D = [-2/3, 2/3]; R = [0, 1]$

B) $D = [-2/3, 2/3]; R = (-\infty, 0]$

C) $D = [-2/3, 2/3]; R = [0, \infty)$

D) $D = (-\infty, -2/3]; R = [0, \infty)$

E) $D = [2/3, \infty); R = [0, 1)$

6. If $f(x) = \frac{2}{3}x + 2$, then $f(x - 3) =$

A) $f(x) - 2$

B) $f(x) + 2$

C) $f(x) - 3$

D) $f(x) + 3$

E) $f(x) + 2/3$

7. If (a, b) is the intersection point of the graphs of $f_1(x) = -3x - 7$ and $f_2(x) = 2x + 13$, then $a + b =$

A) 1

B) -2

C) 4

D) -3

E) 3

8. If $g(x) = 5x^2 - 4x$, then the expression $\frac{g(x+h)-g(x)}{h}$ simplifies to

A) $10x + 5h - 4$

B) $10x + 5h + 4$

C) $10x - 5h + 4$

D) $5x + 5h + 4$

E) $5x - 5h - 4$

9. The domain, in interval notation, of the function $f(x) = \frac{\sqrt{x-2}}{x^2-3x}$ is equal to

A) $[2,3) \cup (3, \infty)$

10. Which one of the following statements is FALSE ?

A) The domain of the function $f(x) = -5$ is $\{-5\}$

B) The range of the relation $x = -7$ is $(-\infty, \infty)$

C) The domain and range of the function $6x - 7y = 0$ are both $(-\infty, \infty)$

D) The slope of a vertical line is undefined

E) The graph of a constant function is a horizontal line

11. If $f(x) = \sqrt{7 - 3x}$, then the Domain D and the Range R , are:

A) D is $(-\infty, \frac{7}{3}]$ and R is $[0, \infty)$

12. If D is the Domain of $y = \frac{5}{x-9}$ and R is the Range of $y = \sqrt{x-1}$

then:

A) $D = (-\infty, 9) \cup (9, \infty)$ and $R = [0, \infty)$

13. If D is the Domain of $y = \frac{1}{\sqrt{x-3}}$ and R is the Range of $y = x^2$ then:

A) $D = (3, \infty)$ and $R = [0, \infty)$

14. The domain of the function $y = \sqrt{\frac{x^2-3x}{2-x}}$ is

A) $(-\infty, 0] \cup (2, 3]$

15. The domain of the function $y = \frac{3}{\sqrt{x-2}}$ is

A) $[0,4) \cup (4, \infty)$

16. The domain of $g(x) = \sqrt{x - x^3}$ is

A) $(-\infty, -1] \cup [0,1]$

17. If $f(x) = \frac{1}{x+1}$, then $\frac{f(1+h)-f(1)}{h}$ is equal to

A) $-\frac{2}{h}$

B) $-\frac{3}{2(2+h)}$

C) $-\frac{1}{2(2+h)}$

D) $-\frac{2}{2+h}$

E) $-2(2+h)$

18. If $f(x) = \sqrt{x}$, then $\frac{f(1+h)-f(1)}{h}$ is equal to

A) $-\frac{1}{1+\sqrt{1+h}}$

B) $\frac{1}{h}$

C) $\frac{1}{\sqrt{1+h}-1}$

D) $\frac{1}{1+\sqrt{1+h}}$

E) $-\frac{1}{h}$

19. Let $h \neq 0$. If $f(x) = x^2 + 5$, then $\frac{f(x) - f(x-h)}{h} =$

A) $2x - h$

B) $2x + h$

C) $2x$

D) $-2x + h$

E) $2x^2 - h$

20. If the domain, in interval notation, of $f(x) = \sqrt{|x - 2| - 1}$ is given by $(-\infty, a] \cup [b, \infty)$, then $a + b =$

A) 4

B) 1

C) 5

D) 3

E) -1

21. If $f(x) = x^2 - 2x + 3$ and $h \neq 0$, then $\frac{f(x+h)-f(x)}{h} =$

A) $2x - h + 2$

B) $2x + h + 2$

C) $x + h - 2$

D) $2x - h - 2$

E) $2x + h - 2$

22. The domain of the function $f(x) = \frac{x-3}{x^3-x^2-9x+9}$ is

A) $(-\infty, -3) \cup (-3, 3) \cup (3, \infty)$

B) $(-\infty, \infty)$

C) $(-\infty, -3) \cup (-3, 1) \cup (1, 3) \cup (3, \infty)$

23. If $f(x) = \frac{1}{x+1}$, then the difference quotient $\frac{f(1)-f(h+1)}{h} =$

A) $\frac{1}{2(h+2)}$

B) $\frac{-1}{2(h+2)}$

C) $\frac{h}{h+2}$

D) $\frac{-1}{h+2}$

E) $\frac{h}{2h+2}$

24. The set of all values of x for which $[\frac{1}{2}x + 1] = -3$, where $[\cdot]$ denotes the greatest integer function, is in the interval

A) $[-8, -6)$

B) $[-4, -3)$

C) $[-7, -5)$

D) $[-3, 0)$

E) $[6, 8)$

25. If $f(x) = \begin{cases} 4x^2 & \text{if } x \leq 0 \\ 2x + 1 & \text{if } 0 < x < 2 \\ |x - 2| & \text{if } x \geq 2 \end{cases}$ then $f(-1) + f(1) + f(5)$

A) 10

26. The relation $y^2 - 1 = x$ is a function if

A) $y < 0$

B) $x < 0$

C) $x > 0$

D) $x > -1$

E) $y > -1$