3.6: Rational Functions and their Graphs

1. If y = 3 is the horizontal asymptote of the function $f(x) = \frac{ax+12}{2x-5}$, then the *x*-intercept of the graph of *f* is

> A) -2 B) $-\frac{1}{2}$ C) $\frac{5}{2}$ D) 3 E) 6

2. If x = 2 is the vertical asymptote of the function $r(x) = \frac{3-(a+1)x}{4-ax}$, then its horizontal asymptote is

A)
$$y = \frac{3}{2}$$

B) $y = -\frac{2}{3}$
C) $y = \frac{3}{4}$
D) $y = 3$
E) $y = 2$

3. The graph of $f(x) = \frac{(2a-1)x+1}{ax-1}$ has the line x = 2 as vertical asymptote, then it has a horizontal asymptote

A) y = 5/3B) y = 2/3C) y = 3/2D) y = 1/3E) y = 2

4. If the graph of the rational function $f(x) = \frac{ax+c}{1-bx}$ is as given below, then the value of a + b + c is equal to



5. If y = -3 is a horizontal asymptote for graph of g(x) =

$$\frac{(Ax^2+x+2)(x+3)}{(2x^2-x-1)(x-3)}$$
, then the vertical asymptote

A) are x = 1 and x = 3B) is x = 1C) are x = 1 and x = -1/2D)are x = 1 and x = 1/2E) is x = 1/2

6. The asymptotes of the graph of the function $f(x) = \frac{x^2 - x - 2}{x^2 + x - 6}$ are

A) one vertical and one horizontal

- B) two vertical and one horizontal
- C) two vertical asymptotes
- D) one horizontal asymptote only
- E) one vertical and two horizontal

7. The rational function $f(x) = \frac{ax^2+2x+1}{bx-4}$ has a horizontal asymptote if

A) $a = 0, b \neq 0$ B) $a \neq 0, b = 0$ C) $a \neq 0, b \neq 0$ D) a = 0, b = 0E) a, b are any real numbers.

8. Which one of the following functions has the graph given below?

A)
$$f(x) = \frac{3-x}{4-x}$$

B) $f(x) = \frac{2-3x}{4-x}$
C) $f(x) = \frac{x-3}{x-4}$
D) $f(x) = \frac{6-3x}{4-x}$
E) $f(x) = \frac{3x-12}{4x-16}$

