

### 5.3: (Trigonometric Functions of Angles)

<p>If <math>\cot \theta = \frac{1}{2}, \pi &lt; \theta &lt; \frac{3\pi}{2}</math>, then <math>\sin \theta + \cos \theta =</math></p> <p>(a) <math>-\frac{3}{\sqrt{5}}</math></p> <p>(b) 3</p> <p>(c) <math>-\frac{1}{\sqrt{5}}</math></p> <p>(d) <math>\frac{3}{\sqrt{5}}</math></p> <p>(e) <math>\frac{1}{\sqrt{5}}</math></p>	<p>Trigonometric Functions of Angles.</p>
<p>If <math>\tan x = \frac{12}{5}</math> for all <math>x</math> is in the third quadrant, then <math>\cos x =</math></p> <p>(a) <math>-\frac{5}{13}</math></p> <p>(b) <math>\frac{5}{13}</math></p> <p>(c) <math>-\frac{12}{13}</math></p> <p>(d) <math>\frac{12}{13}</math></p> <p>(e) <math>\frac{13}{5}</math></p>	<p>Trigonometric Functions of Angles.</p>
<p>If <math>\cos \theta = -\frac{1}{2}</math> and <math>\sin \theta &gt; 0</math>, then <math>\cot \theta + \csc \theta =</math></p> <p>A) <math>\frac{\sqrt{3}}{3}</math></p> <p>B) <math>\frac{1}{2}</math></p> <p>C) <math>-\frac{\sqrt{3}}{2}</math></p> <p>D) <math>-\sqrt{3}</math></p> <p>E) <math>\sqrt{3}</math></p>	<p>Trigonometric Functions of Angles.</p>

<p>If <math>\sec \theta = -5</math> and <math>\sin \theta &gt; 0</math>, then <math>\tan \theta - \sin \theta =</math></p> <p>A) <math>\frac{8\sqrt{6}}{5}</math></p> <p>B) <math>12\sqrt{6}</math></p> <p>C) <math>-\frac{12\sqrt{6}}{5}</math></p> <p>D) <math>-2\sqrt{6}</math></p> <p>E) <math>-\frac{8\sqrt{6}}{5}</math></p>	<p>Trigonometric Functions of Angles.</p>
<p>If <math>\theta</math> is in quadrant IV and <math>\sec \theta = \frac{x+4}{x}</math>, where <math>x &gt; 0</math>, then <math>\tan \theta =</math></p> <p>A) <math>-\frac{2\sqrt{2x+4}}{x}</math></p> <p>B) <math>-\frac{\sqrt{2x+4}}{2x}</math></p> <p>C) <math>\frac{2\sqrt{x+1}}{x}</math></p> <p>D) <math>-\frac{4\sqrt{x+4}}{x}</math></p> <p>E) <math>\frac{4\sqrt{x+4}}{x}</math></p>	<p>Trigonometric Functions of Angles.</p>
<p>If <math>x</math> is in the third quadrant, then <math>\cot x</math> in terms of <math>\sin x</math> is</p> <p>(a) <math>-\frac{\sqrt{1-\sin^2 x}}{\sin x}</math></p> <p>(b) <math>\frac{\sqrt{1-\sin^2 x}}{\sin x}</math></p> <p>(c) <math>-\frac{\sin x}{\sqrt{1+\sin^2 x}}</math></p> <p>(d) <math>\frac{\sin x}{\sqrt{1+\sin^2 x}}</math></p> <p>(e) <math>-\frac{\sqrt{1-\sin x}}{\sin x}</math></p>	<p>Trigonometric Functions of Angles.</p>

<p>If <math>x</math> is in the third quadrant, then <math>\cot x</math> in terms of <math>\sec x</math> is</p> <p>A) <math>\frac{\sqrt{\sec^2 x - 1}}{\sec^2 x - 1}</math></p> <p>B) <math>-\frac{\sqrt{\sec^2 x - 1}}{\sec^2 x - 1}</math></p> <p>C) <math>-\frac{\sqrt{\sec^2 x + 1}}{\sec^2 x - 1}</math></p> <p>D) <math>\frac{\sqrt{\sec^2 x - 1}}{\sec^2 x + 1}</math></p> <p>E) <math>-\frac{1}{\sec^2 x - 1}</math></p>	<p>Trigonometric Functions of Angles.</p>
<p>If <math>\csc \theta = \frac{x+1}{x}</math>, <math>x &gt; 0</math>, then <math>\cot \theta =</math></p> <p>A) <math>\frac{\sqrt{1+2x}}{x}</math></p> <p>B) <math>\frac{\sqrt{2x-1}}{x}</math></p> <p>C) <math>\frac{\sqrt{x^2+2x}}{x}</math></p> <p>D) <math>\frac{\sqrt{2x^2+2x+1}}{x}</math></p> <p>E) <math>\frac{1}{x}</math></p>	<p>Trigonometric Functions of Angles.</p>
<p>If <math>\cot \theta = u</math> and <math>\theta</math> is in the third quadrant, then <math>\cot \theta \sec \theta =</math></p> <p>A) <math>-\sqrt{1+u^2}</math></p> <p>B) <math>\sqrt{1+u^2}</math></p> <p>C) <math>\sqrt{1-u^2}</math></p> <p>D) <math>-\sqrt{1-u^2}</math></p> <p>E) <math>\sqrt{u^2-1}</math></p>	<p>Trigonometric Functions of Angles.</p>

<p>If <math>\cot \theta = m</math>, where <math>\pi &lt; \theta &lt; \frac{3\pi}{2}</math>, then <math>\cos \theta</math> is equal to</p> <p>A) <math>-\frac{m\sqrt{1+m^2}}{1+m^2}</math></p> <p>B) <math>-\frac{m\sqrt{1-m^2}}{1-m^2}</math></p> <p>C) <math>\frac{-\sqrt{1-m^2}}{1-m^2}</math></p> <p>D) <math>\frac{-\sqrt{1+m^2}}{1+m^2}</math></p> <p>E) <math>\sqrt{1-m^2}</math></p>	<p>Trigonometric Functions of Angles.</p>
<p>If <math>\cos \theta = -\frac{2}{3}</math>, <math>\sin \theta &lt; 0</math>, then <math>\csc \theta + \tan \theta</math></p> <p>A) <math>-\frac{\sqrt{5}}{10}</math></p> <p>B) <math>\frac{11\sqrt{5}}{10}</math></p> <p>C) <math>-\frac{\sqrt{13}}{10}</math></p> <p>D) <math>\frac{3\sqrt{5}}{10}</math></p> <p>E) <math>-\frac{7\sqrt{5}}{10}</math></p>	<p>Trigonometric Functions of Angles.</p>
<p>If <math>\tan \theta = \frac{3}{4}</math>, where <math>\theta</math> is in the third quadrant, then <math>\csc \theta =</math></p> <p>A) <math>-\frac{5}{3}</math></p> <p>B) <math>\frac{5}{3}</math></p> <p>C) <math>-\frac{13}{5}</math></p> <p>D) <math>-\frac{5}{4}</math></p> <p>E) <math>\frac{5}{4}</math></p>	<p>Trigonometric Functions of Angles.</p>

<p>Which one of the following statements is possible?</p> <p>A) <math>\tan \theta = -\frac{\sqrt{3}}{2}</math> and <math>\sec \theta = \frac{\sqrt{7}}{2}</math></p> <p>B) <math>\sin \theta = \frac{\pi}{2}</math></p> <p>C) <math>\csc \theta = -\frac{1}{2}</math> and <math>\sin \theta = -2</math></p> <p>D) <math>\cos \theta = -\frac{3}{2}</math> and <math>\sec \theta = -\frac{2}{3}</math></p> <p>E) <math>\sec \theta = 0</math></p>	<p>Trigonometric Functions of Angles.</p>
<p>If <math>\cot \theta = \frac{1}{2}</math> where <math>\pi &lt; \theta &lt; \frac{3\pi}{2}</math>, then <math>\sin \theta - \cos \theta =</math></p> <p>A) <math>-\frac{\sqrt{5}}{5}</math></p> <p>B) <math>-\frac{2\sqrt{5}}{5}</math></p> <p>C) <math>\frac{3\sqrt{5}}{5}</math></p> <p>D) <math>\frac{2\sqrt{5}}{5}</math></p> <p>E) <math>-\frac{3\sqrt{5}}{5}</math></p>	<p>Trigonometric Functions of Angles.</p>
<p>Which one of the following statements is FALSE for any angle <math>\alpha</math> in the domain of the functions?</p> <p>A) <math>\sin \alpha + \cos \alpha = 1</math></p> <p>B) <math>-1 \leq \sin \alpha \leq 1</math></p> <p>C) <math>1 \leq  \sec \alpha </math></p> <p>D) <math>1 \leq  \csc \alpha </math></p> <p>E) <math>-\infty &lt; \tan \alpha &lt; \infty</math></p>	<p>Trigonometric Functions of Angles.</p>

<p>If <math>\tan \theta = -\frac{5}{3}</math> and <math>\theta</math> is in the second quadrant, then <math>\frac{\csc \theta - \cot \theta}{\cos \theta} =</math></p> <p>A) <math>-\frac{34+3\sqrt{34}}{15}</math></p> <p>B) <math>\frac{3+\sqrt{34}}{15}</math></p> <p>C) <math>-\frac{\sqrt{34}}{15}</math></p> <p>D) <math>\frac{3\sqrt{34}-34}{34}</math></p> <p>E) <math>\frac{34}{9} - \frac{\sqrt{34}}{5}</math></p>	<p>Trigonometric Functions of Angles.</p>
<p>The exact value of <math>-\tan(780^\circ)\sin(570^\circ) - \sec(-585^\circ)</math> is</p> <p>A) <math>\frac{\sqrt{3}+2\sqrt{2}}{2}</math></p> <p>B) <math>\frac{\sqrt{3}-2\sqrt{2}}{2}</math></p> <p>C) <math>\frac{2\sqrt{3}-3\sqrt{2}}{2}</math></p> <p>D) <math>\frac{\sqrt{3}-\sqrt{2}}{2}</math></p> <p>E) <math>\frac{\sqrt{3}+\sqrt{2}}{2}</math></p>	<p>Trigonometric Functions of Angles.</p>
<p>Which one of the following statements is TRUE?</p> <p>A) If <math>\tan \theta = \sqrt{3}</math> and <math>\theta</math> is in Quadrant III, then <math>\cos \theta = -\frac{1}{2}</math>.</p> <p>B) If <math>\cot \theta = 2</math>, then <math>\sin \theta = 2</math> and <math>\cos \theta = 1</math>.</p> <p>C) If <math>\sec \theta &gt; 0</math> and <math>\csc \theta &gt; 0</math>, then <math>\theta</math> lies in Quadrant II.</p> <p>D) If <math>90^\circ &lt; \theta &lt; 180^\circ</math>, then <math>\sin(2\theta)</math> is positive.</p> <p>E) If <math>\sec \theta = \frac{10}{3}</math>, then <math>\sin \theta = \frac{3}{10}</math>.</p>	<p>Trigonometric Functions of Angles.</p>

<p>If <math>\cot \theta = \frac{2}{\sqrt{5}}</math>, <math>\sec \theta &lt; 0</math>, then <math>\sin \theta \cos \theta =</math></p> <p>A) <math>\frac{2\sqrt{5}}{9}</math></p> <p>B) <math>2\sqrt{5}</math></p> <p>C) <math>-\frac{2\sqrt{5}}{9}</math></p> <p>D) <math>\frac{\sqrt{5}}{3}</math></p> <p>E) <math>-\frac{\sqrt{5}}{9}</math></p>	<p>Trigonometric Functions of Angles.</p>
<p><math>\sec\left(-\frac{23\pi}{6}\right) \cot\left(\frac{16\pi}{3}\right) =</math></p> <p>A) <math>\frac{2}{3}</math></p> <p>B) <math>-\frac{2}{3}</math></p> <p>C) 2</p> <p>D) <math>-\frac{3}{2}</math></p> <p>E) -3</p>	<p>Trigonometric Functions of Angles.</p>
<p>The exact value of <math>\tan\left(-\frac{7\pi}{6}\right) + \sec\left(-\frac{\pi}{6}\right)</math> is equal to</p> <p>(a) <math>\frac{\sqrt{3}}{3}</math></p> <p>(b) <math>-\frac{\sqrt{3}}{3}</math></p> <p>(c) <math>-\frac{3\sqrt{3}}{2}</math></p> <p>(d) <math>-\frac{2\sqrt{3}}{3}</math></p> <p>(e) <math>\frac{\sqrt{3}}{2}</math></p>	<p>Trigonometric Functions of Angles.</p>

<p>The value of <math>\cot\left(-\frac{17\pi}{3}\right) + \sin\left(\frac{11\pi}{6}\right) =</math></p> <p>A) <math>\frac{2\sqrt{3}-3}{6}</math></p> <p>B) <math>\frac{2\sqrt{3}+1}{3}</math></p> <p>C) <math>\frac{\sqrt{3}-3}{3}</math></p> <p>D) <math>\frac{2\sqrt{3}-1}{6}</math></p> <p>E) <math>\frac{\sqrt{3}+2}{2}</math></p>	<p>Trigonometric Functions of Angles.</p>
<p><math>\tan(570^\circ) + \csc(-1020^\circ) =</math></p> <p>A) <math>\sqrt{3}</math></p> <p>B) <math>-\sqrt{3}</math></p> <p>C) <math>-\frac{\sqrt{3}}{3}</math></p> <p>D) <math>\frac{\sqrt{3}}{3}</math></p> <p>E) <math>\sqrt{3} + 2</math></p>	<p>Trigonometric Functions of Angles.</p>
<p>If <math>u = \sin 780^\circ</math> and <math>v = \cot(-950^\circ) + \tan 220^\circ</math>, then <math>4(u^2 + v) =</math></p> <p>A) 3</p> <p>B) <math>\sqrt{3}</math></p> <p>C) <math>3 + 4\cot 40^\circ</math></p> <p>D) <math>\sqrt{3} + 4\tan 40^\circ</math></p> <p>E) -1</p>	<p>Trigonometric Functions of Angles.</p>



<p>If <math>\tan \theta = -\frac{2\sqrt{5}}{5}</math> and <math>\sec \theta = -\frac{3\sqrt{5}}{5}</math> then, <math>12\csc \theta =</math></p> <p>A) 18  B) 8  C) -18  D) -8  E) <math>-3\sqrt{5}</math></p>	<p>Trigonometric Functions of Angles.</p>
<p><math>\sqrt{3}\tan(750^\circ) + 2\sec(-300^\circ) =</math></p> <p>A) 5  B) -3  C) 7  D) -1  E) 2</p>	<p>Trigonometric Functions of Angles.</p>
<p>If <math>\cos 160^\circ = A</math>, then <math>\cos 340^\circ + \sec 200^\circ</math> equals to</p> <p>A) <math>\frac{1-A^2}{A}</math>  B) <math>\frac{1+A^2}{A}</math>  C) <math>\frac{A^2-1}{A}</math>  D) <math>\frac{1}{A}</math>  E) <math>A^2 + 1</math></p>	<p>Trigonometric Functions of Angles.</p>

<p>The exact value of <math>\sec(-480^\circ) - \cot\frac{3\pi}{4}</math> is</p> <p>A) - 1</p> <p>B) -3</p> <p>C) <math>\frac{3-2\sqrt{3}}{3}</math></p> <p>D) 3</p> <p>E) <math>\frac{3+2\sqrt{3}}{3}</math></p>	<p>Trigonometric Functions of Angles.</p>
<p><math>\sec\frac{25\pi}{6} - \tan(-510^\circ) =</math></p> <p>A) <math>\frac{\sqrt{3}}{3}</math></p> <p>B) <math>\frac{2\sqrt{3}}{3}</math></p> <p>C) <math>\sqrt{3}</math></p> <p>D) <math>-\frac{\sqrt{3}}{3}</math></p> <p>E) <math>-\frac{2\sqrt{3}}{3}</math></p>	<p>Trigonometric Functions of Angles.</p>
<p><math>\tan\left(\frac{23\pi}{6}\right) + \csc\left(\frac{11\pi}{6}\right) =</math></p> <p>A) <math>\frac{-\sqrt{3}-6}{3}</math></p> <p>B) <math>\frac{-\sqrt{3}+6}{3}</math></p> <p>C) <math>\frac{-2\sqrt{3}-3}{6}</math></p> <p>D) <math>\frac{\sqrt{3}+2}{3}</math></p> <p>E) <math>\frac{\sqrt{3}-2}{3}</math></p>	<p>Trigonometric Functions of Angles.</p>

<p>If <math>\alpha</math> is the reference angle of <math>-30^\circ</math> and <math>\beta</math> is the smallest positive coterminal angle of <math>-670^\circ</math>, then <math>\alpha + \beta =</math></p> <p>A) <math>80^\circ</math>  B) <math>380^\circ</math>  C) <math>110^\circ</math>  D) <math>200^\circ</math>  E) <math>20^\circ</math></p>	<p>Trigonometric Functions of Angles.</p>
<p>If <math>\alpha = 475^\circ</math> and <math>\beta = -\frac{11\pi}{6}</math> are two angles in standard position, <math>2\alpha + \beta</math> is in the</p> <p>A) third quadrant  B) first quadrant  C) second quadrant  D) fourth quadrant  E) quadrantal angle</p>	<p>Trigonometric Functions of Angles.</p>
<p>If <math>\pi &lt; \theta &lt; \frac{3\pi}{2}</math> and <math>\cot \theta = \frac{3\sqrt{7}}{7}</math>, then <math>\cos \theta =</math></p> <p>A) <math>-\frac{3}{4}</math>  B) <math>-\frac{4}{3}</math>  C) <math>\frac{\sqrt{7}}{4}</math>  D) <math>-\frac{\sqrt{7}}{3}</math>  E) <math>\frac{3}{4}</math></p>	<p>Trigonometric Functions of Angles.</p>

<p>The exact value of <math>\tan(675^\circ)\cos(-240^\circ) - \csc(495^\circ)</math> is</p> <p>A) <math>\frac{1-2\sqrt{2}}{2}</math></p> <p>B) <math>\frac{1+2\sqrt{2}}{2}</math></p> <p>C) <math>\frac{-1-2\sqrt{2}}{2}</math></p> <p>D) <math>\frac{1-\sqrt{2}}{2}</math></p> <p>E) <math>\frac{1+\sqrt{2}}{2}</math></p>	<p>Trigonometric Functions of Angles.</p>
<p><math>\tan\left(-\frac{5\pi}{3}\right) + \csc\left(\frac{23\pi}{6}\right) =</math></p> <p>A) <math>\sqrt{3} - 2</math></p> <p>B) <math>\sqrt{3} + 2</math></p> <p>C) <math>2 - \sqrt{3}</math></p> <p>D) <math>\frac{\sqrt{3}+2}{2}</math></p> <p>E) <math>\frac{\sqrt{3}-2}{2}</math></p>	<p>Trigonometric Functions of Angles.</p>
<p>The exact value of <math>\sec\left(-\frac{19\pi}{4}\right) \cdot \tan\left(\frac{17\pi}{3}\right) + \csc\left(\frac{11\pi}{6}\right)</math> is equal to</p> <p>A) <math>\sqrt{6} - 2</math></p> <p>B) <math>\frac{\sqrt{6}-6}{3}</math></p> <p>C) <math>\frac{2\sqrt{6}-1}{2}</math></p> <p>D) <math>-\sqrt{6} + 2</math></p> <p>E) <math>-\sqrt{6} - 2</math></p>	<p>Trigonometric Functions of Angles.</p>

<p>Which one of the following statements is TRUE for <math>-90^\circ &lt; \theta &lt; 90^\circ</math>.</p> <p>A) <math>\sec(\theta - 180^\circ)</math> is negative</p> <p>B) <math>\cos \frac{\theta}{2}</math> is negative</p> <p>C) <math>\cos(\theta + 180^\circ)</math> is positive</p> <p>D) <math>\sin(\theta - 90^\circ)</math> is positive</p> <p>E) <math>\sec(-\theta)</math> is negative</p>	<p>Trigonometric Functions of Angles.</p>
<p><math>\sec(480^\circ) =</math></p> <p>A) <math>-\frac{2\sqrt{3}}{3}</math></p> <p>B) 2</p> <p>C) <math>\frac{2\sqrt{3}}{3}</math></p> <p>D) -2</p> <p>E) -1</p>	<p>Trigonometric Functions of Angles.</p>
<p><math>4\sin(-870^\circ) + \tan 143^\circ + \cot 53^\circ =</math></p> <p>A) -2</p> <p>B) 2</p> <p>C) <math>2\sqrt{3}</math></p> <p>D) <math>-2\sqrt{3}</math></p> <p>E) <math>2\sqrt{2}</math></p>	<p>Trigonometric Functions of Angles.</p>

<p><math>4\sin(-510^\circ)\cos 300^\circ + \cot 199^\circ - \tan 251^\circ =</math></p> <p>A) -1</p> <p>B) 1</p> <p>C) <math>1 - 2\sqrt{3}</math></p> <p>D) <math>-1 + 2\sqrt{3}</math></p> <p>E) <math>-1 - 2\sqrt{3}</math></p>	<p>Trigonometric Functions of Angles.</p>
<p>Which one of the following statements is TRUE?</p> <p>A) <math>\sec(-89^\circ) &gt; 0</math></p> <p>B) <math>\cot(-100^\circ) &lt; 0</math></p> <p>C) <math>\cos 178^\circ &gt; 0</math></p> <p>D) <math>\tan 340^\circ &gt; 0</math></p> <p>E) <math>\sin 370^\circ &lt; 0</math></p>	<p>Trigonometric Functions of Angles.</p>
<p>Which one of the following statements is FALSE?</p> <p>A) <math>(\sin \theta + \cos \theta)^2 = 1</math> for all angles <math>\theta</math>.</p> <p>B) If <math>90^\circ &lt; \theta &lt; 180^\circ</math>, then <math>\cot\left(\frac{\theta}{2}\right)</math> is positive</p> <p>C) If <math>\sec \theta &lt; 0</math> and <math>\csc \theta &lt; 0</math>, then <math>\theta</math> lies in Quadrant III.</p> <p>D) If <math>\tan \theta &lt; 0</math> and <math>\cot \theta &lt; 0</math>, then <math>\theta</math> lies in Quadrant II or IV.</p> <p>E) If <math>\csc \theta = 2</math> and <math>\theta</math> in quadrant II, then <math>\cos \theta = -\frac{\sqrt{3}}{2}</math></p>	<p>Trigonometric Functions of Angles.</p>

$\cos\left(\frac{7\pi}{4}\right)\tan\left(\frac{4\pi}{3}\right) + \cos\left(\frac{7\pi}{6}\right) =$ <p>A) <math>\frac{\sqrt{6}-\sqrt{3}}{2}</math></p> <p>B) <math>\frac{\sqrt{6}+\sqrt{3}}{2}</math></p> <p>C) <math>\frac{\sqrt{2}+\sqrt{3}}{2}</math></p> <p>D) <math>\frac{\sqrt{2}-\sqrt{3}}{2}</math></p> <p>E) <math>\frac{\sqrt{6}-\sqrt{3}}{4}</math></p>	<p>Trigonometric Functions of Angles.</p>
<p>The exact value of <math>\sec(-480^\circ) + \csc\left(\frac{71\pi}{6}\right)</math> is</p> <p>A) -4</p> <p>B) <math>-\frac{1}{2}</math></p> <p>C) -2</p> <p>D) <math>-\frac{1}{4}</math></p> <p>E) -3</p>	<p>Trigonometric Functions of Angles.</p>
<p>The value of <math>\sin 150^\circ + \tan \frac{5\pi}{4} + \sec 300^\circ</math> is</p> <p>A) <math>\frac{7}{2}</math></p> <p>B) <math>\frac{3}{2}</math></p> <p>C) <math>\frac{5-\sqrt{2}}{2}</math></p> <p>D) <math>\frac{1}{2}</math></p>	<p>Trigonometric Functions of Angles.</p>

<p>The exact value of <math>\sin(-210^\circ) + \cot(735^\circ) + \tan(285^\circ)</math> is</p> <p>A) <math>\frac{1}{2}</math></p> <p>B) <math>\frac{\sqrt{2}}{2}</math></p> <p>C) <math>-\frac{\sqrt{3}}{2}</math></p> <p>D) <math>-\frac{1}{2}</math></p> <p>E) <math>\frac{\sqrt{3}}{2}</math></p>	<p>Trigonometric Functions of Angles.</p>
<p>If <math>2\sin \theta = -3\cos \theta, \frac{3\pi}{2} &lt; \theta &lt; 2\pi</math>, then <math>\sin \theta - \cos \theta =</math></p> <p>A) <math>-\frac{5}{\sqrt{13}}</math></p> <p>B) <math>\frac{5}{\sqrt{13}}</math></p> <p>C) <math>-\frac{1}{\sqrt{13}}</math></p> <p>D) <math>\frac{1}{\sqrt{13}}</math></p> <p>E) -1</p>	<p>Trigonometric Functions of Angles.</p>
<p>The value of <math>\cos(-510^\circ)\csc(300^\circ) + \tan\left(-\frac{9\pi}{4}\right)</math> is</p> <p>(a) 0</p> <p>(b) <math>\sqrt{3} - 1</math></p> <p>(c) -2</p> <p>(d) <math>\frac{4}{3}</math></p> <p>(e) <math>\frac{3}{4}</math></p>	<p>Trigonometric Functions of Angles.</p>



<p><math>4\sin(-510^\circ)\cos 300^\circ + \cot 199^\circ - \tan 251^\circ =</math></p> <p>A) -1</p> <p>B) 1</p> <p>C) <math>1 - 2\sqrt{3}</math></p> <p>D) <math>-1 + 2\sqrt{3}</math></p> <p>E) <math>-1 - 2\sqrt{3}</math></p>	<p>Trigonometric Functions of Angles.</p>
<p>If <math>\tan 20^\circ = a</math>, then <math>\tan 160^\circ + \tan(-380^\circ) =</math></p> <p>A) <math>-2a</math></p> <p>B) 0</p> <p>C) <math>2a</math></p> <p>D) <math>\sqrt{1 + a^2}</math></p> <p>E) <math>\frac{1-a}{a}</math></p>	<p>Trigonometric Functions of Angles.</p>
<p>If <math>\tan 324^\circ = x</math>, then <math>\csc 36^\circ =</math></p> <p>A) <math>-\frac{\sqrt{x^2+1}}{x}</math></p> <p>B) <math>\frac{\sqrt{x^2+1}}{x}</math></p> <p>C) <math>\sqrt{x^2 + 1}</math></p> <p>D) <math>-\sqrt{x^2 + 1}</math></p> <p>E) <math>\frac{1}{x}</math></p>	<p>Trigonometric Functions of Angles.</p>

<p>If <math>\sec \frac{9\pi}{5} = x</math>, then <math>\tan \frac{\pi}{5} =</math></p> <p>A) <math>\sqrt{x-1}</math>  B) <math>\frac{\sqrt{x^2-1}}{x}</math>  C) <math>\sqrt{x^2-1}</math>  D) <math>\sqrt{x+1}</math>  E) <math>\sqrt{x^2+1}</math></p>	<p>Trigonometric Functions of Angles.</p>
<p>For any angle <math>\theta</math>, which one of the following is not possible?</p> <p>A) <math>\cos \theta = -\frac{4}{3}</math> and <math>\sec \theta = -\frac{3}{4}</math>  B) <math>\sin^2(-\theta) + \cos^2(-\theta) = 1</math>  C) <math>\tan \theta = 4</math> and <math>\cot \theta = \frac{1}{4}</math>  D) <math>\cot^2 \theta = \csc^2 \theta - 1</math>  E) <math>\csc \theta = -5</math></p>	<p>Trigonometric Functions of Angles.</p>
<p>The exact value of <math>\sec \frac{23\pi}{6} \cot \frac{13\pi}{3} - \sin \frac{7\pi}{4}</math> is equal to</p> <p>A) <math>\frac{4+3\sqrt{2}}{6}</math>  B) <math>\frac{4-3\sqrt{2}}{6}</math>  C) <math>-\frac{4+3\sqrt{2}}{6}</math>  D) <math>\frac{3\sqrt{2}-4}{6}</math>  E) <math>\frac{4+\sqrt{2}}{2}</math></p>	<p>Trigonometric Functions of Angles.</p>

$\sin\left(\frac{7\pi}{4}\right) \tan 600^\circ + \cos\left(-\frac{7\pi}{6}\right) =$ <p>A) <math>-\frac{\sqrt{3}}{2}(\sqrt{2} + 1)</math></p> <p>B) <math>\frac{\sqrt{3}}{2}(\sqrt{2} - 1)</math></p> <p>C) <math>\frac{\sqrt{3}}{2}(1 - \sqrt{2})</math></p> <p>D) <math>\frac{\sqrt{3}}{2}(\sqrt{2} + 1)</math></p> <p>E) <math>-\frac{\sqrt{2}}{2}(\sqrt{3} + 1)</math></p>	<p>Trigonometric Functions of Angles.</p>
<p>If <math>-90^\circ &lt; \theta &lt; 90^\circ</math>, then</p> <p>A) <math>\sin(\theta + 90^\circ) &gt; 0</math> and <math>\sec\frac{\theta}{2} &gt; 0</math></p> <p>B) <math>\sin(\theta + 90^\circ) &lt; 0</math> and <math>\sec\frac{\theta}{2} &gt; 0</math></p> <p>C) <math>\sin(\theta + 90^\circ) &gt; 0</math> and <math>\sec\frac{\theta}{2} &lt; 0</math></p> <p>D) <math>\sin(\theta + 90^\circ) &lt; 0</math> and <math>\sec\frac{\theta}{2} &lt; 0</math></p> <p>E) <math>\tan \theta &lt; 0</math> and <math>\cos \theta &gt; 0</math></p>	<p>Trigonometric Functions of Angles.</p>
<p>The exact value of <math>\csc(225^\circ) \cdot \tan(-240^\circ) + \sin 150^\circ</math> is</p> <p>(a) <math>\frac{1}{2} + \sqrt{6}</math></p> <p>(b) <math>\sqrt{3} + \frac{1}{2}</math></p> <p>(c) <math>\frac{\sqrt{2}+2\sqrt{3}}{\sqrt{3}}</math></p> <p>(d) <math>\frac{\sqrt{6}+4\sqrt{3}}{\sqrt{2}}</math></p> <p>(e) <math>\sqrt{6} + 2\sqrt{3}</math></p>	<p>Trigonometric Functions of Angles.</p>

<p>If <math>\tan(71^\circ) = b</math>, then <math>\csc^2(19^\circ) + 1 =</math></p> <p>A) <math>b^2 + 2</math></p> <p>B) <math>b^2 + 1</math></p> <p>C) 1</p> <p>D) <math>b^2</math></p> <p>E) <math>b^2 - 1</math></p>	<p>Trigonometric Functions of Angles.</p>
<p><math>\cos \frac{17\pi}{4} - \tan 765^\circ \csc \frac{11\pi}{6} =</math></p> <p>A) <math>\frac{\sqrt{2}+4}{2}</math></p> <p>B) <math>4\sqrt{2}</math></p> <p>C) <math>\frac{\sqrt{2}-4}{4}</math></p> <p>D) <math>\frac{\sqrt{3}+1}{2}</math></p> <p>E) <math>\frac{2\sqrt{2}}{3}</math></p>	<p>Trigonometric Functions of Angles.</p>
<p><math>\sin\left(-\frac{7\pi}{4}\right) + \tan(870^\circ)</math></p> <p>A) <math>\frac{3\sqrt{2}-2\sqrt{3}}{6}</math></p> <p>B) <math>\frac{3}{2}</math></p> <p>C) <math>\frac{3\sqrt{2}+2\sqrt{3}}{6}</math></p> <p>D) <math>\frac{2\sqrt{2}-3\sqrt{3}}{6}</math></p> <p>E) <math>\frac{-3\sqrt{2}-2\sqrt{3}}{6}</math></p>	<p>Trigonometric Functions of Angles.</p>

<p>If <math>\tan(24^\circ) = t</math> then <math>\tan(516^\circ) + \cot(156^\circ) =</math></p> <p>A) <math>\frac{-t^2-1}{t}</math></p> <p>B) <math>\frac{t^2-1}{t}</math></p> <p>C) <math>\frac{t^2+1}{t}</math></p> <p>D) <math>\frac{-t-1}{t}</math></p> <p>E) <math>\frac{t+1}{t}</math></p>	<p>Trigonometric Functions of Angles.</p>
<p>Which one of the following statements is TRUE?</p> <p>(a) If <math>90^\circ &lt; \theta &lt; 180^\circ</math> then, <math>\sin(2\theta)</math> is negative.</p> <p>(b) <math>\sin \theta + \cos \theta = 1</math> for all <math>\theta</math>.</p> <p>(c) If <math>\cot \theta = \frac{1}{2}</math>, then <math>\sin \theta = 1</math> and <math>\cos \theta = 2</math>.</p> <p>(d) <math>\sec \theta = -0.3</math>, for some <math>\theta</math> where <math>\frac{\pi}{2} &lt; \theta &lt; \pi</math>.</p> <p>(e) If <math>\sec \theta &gt; 0</math>, <math>\csc \theta &gt; 0</math>, then <math>\theta</math> is in the second quadrant.</p>	<p>Trigonometric Functions of Angles.</p>
<p>Which one of the following statements is FALSE?</p> <p>A) If <math>90^\circ \leq \theta \leq 180^\circ</math> then <math>\tan\left(\frac{\theta}{2}\right)</math> is negative.</p> <p>B) If <math>\tan \theta = \frac{1}{2}</math> then the terminal side of <math>\theta</math> lies in quadrant I or quadrant III .</p> <p>C) The range of <math>\tan \theta</math> is <math>(-\infty, \infty)</math>.</p> <p>D) If <math>0 \leq \theta &lt; \frac{\pi}{2}</math>, then <math>\tan^2 \theta = \sec^2 \theta - 1</math>.</p> <p>E) If <math>\tan(-15^\circ) = \beta</math> then, <math>\tan(15^\circ) = -\beta</math>.</p>	<p>Trigonometric Functions of Angles.</p>

<p>Which one of the following statements is FALSE?</p> <p>A) If <math>0 \leq \theta &lt; \frac{\pi}{2}</math> and <math>\tan \theta = \frac{1}{2}</math>, then <math>\sin \theta = 1</math> and <math>\cos \theta = 2</math></p> <p>B) If <math>0 \leq \theta &lt; \frac{\pi}{2}</math>, then <math>\sec^2 \theta - \tan^2 \theta = 1</math></p> <p>C) If <math>0 \leq \theta &lt; \frac{\pi}{2}</math>, then <math>\sin\left(\frac{\theta}{2}\right)</math> is positive</p> <p>D) The range of <math>\tan \theta</math> is <math>(-\infty, \infty)</math></p> <p>E) <math>\sin\left(-\frac{\pi}{3}\right) = \sin\left(\frac{5\pi}{3}\right)</math></p>	<p>Trigonometric Functions of Angles.</p>
<p>Decide which one of the following statements is possible.</p> <p>A) <math>\cot \theta = 0.93</math></p> <p>B) <math>\cos \theta = -\frac{4}{3}</math></p> <p>C) <math>\tan \theta = \frac{3}{2}</math> and <math>\cot \theta = -\frac{3}{2}</math></p> <p>D) <math>\csc \theta = -\frac{1}{2}</math> and <math>\sin \theta = -2</math></p> <p>E) <math>\sec \theta = -0.3</math></p>	<p>Trigonometric Functions of Angles.</p>
<p><math>\tan(420^\circ) + \sec(495^\circ)\csc(225^\circ) =</math></p> <p>A) <math>\sqrt{3} + 2</math></p> <p>B) <math>-\sqrt{3} + 2</math></p> <p>C) <math>-\sqrt{3} - 2</math></p> <p>D) <math>\frac{\sqrt{3}}{3} + 2</math></p> <p>E) <math>-\frac{\sqrt{3}}{3} + \frac{1}{2}</math></p>	<p>Trigonometric Functions of Angles.</p>

<p>If <math>\alpha</math> is the least positive coterminal angle with the angle <math>\frac{65\pi}{9}</math>, and <math>\beta</math> is the reference angle of the angle <math>\frac{5\pi}{9}</math>, then <math>\alpha + \beta =</math></p> <p>A) <math>\frac{5\pi}{3}</math></p> <p>B) <math>\frac{11\pi}{9}</math></p> <p>C) <math>\frac{\pi}{9}</math></p> <p>D) <math>\frac{-\pi}{9}</math></p> <p>E) <math>\frac{13\pi}{9}</math></p>	<p>Reference &amp; Co-terminal Angles.</p>
<p>The reference angle of <math>\theta = \frac{11\pi}{15}</math>, in degrees, is equal to</p> <p>A) <math>48^\circ</math></p> <p>B) <math>32^\circ</math></p> <p>C) <math>49^\circ</math></p> <p>D) <math>38^\circ</math></p> <p>E) <math>35^\circ</math></p>	<p>Reference &amp; Co-terminal Angles.</p>
<p>If R is the reference angle of <math>1945^\circ</math> and Q is the smallest positive coterminal angle of <math>-950^\circ</math>, then <math>R + Q</math></p> <p>A) <math>165^\circ</math></p> <p>B) <math>155^\circ</math></p> <p>C) <math>175^\circ</math></p> <p>D) <math>275^\circ</math></p> <p>E) <math>255^\circ</math></p>	<p>Reference &amp; Co-terminal Angles.</p>

<p>If the reference angle of 10 radians is <math>10 - n\pi</math>, then <math>n =</math></p> <p>A) 3</p> <p>B) 6</p> <p>C) 4</p> <p>D) 7</p> <p>E) 5</p>	<p>Reference &amp; Co-terminal Angles.</p>
<p>The reference angle of the angle <math>\theta = \frac{25\pi}{7}</math> is equal to</p> <p>A) <math>\frac{3\pi}{7}</math></p> <p>B) <math>\frac{2\pi}{7}</math></p> <p>C) <math>\frac{5\pi}{7}</math></p> <p>D) <math>\frac{\pi}{7}</math></p> <p>E) <math>\frac{4\pi}{7}</math></p>	<p>Reference &amp; Co-terminal Angles.</p>
<p>The smallest positive coterminal angle of <math>\theta = \frac{23\pi}{7}</math> is</p> <p>A) in the third quadrant</p> <p>B) in the first quadrant</p> <p>C) in the fourth quadrant</p> <p>D) a quadrantal angle</p> <p>E) in the second quadrant</p>	<p>Reference &amp; Co-terminal Angles.</p>



<p>The sum of all coterminal angles with <math>\frac{2\pi}{3}</math> between <math>2\pi</math> and <math>6\pi</math> is</p> <p>A) <math>\frac{22\pi}{3}</math></p> <p>B) <math>\frac{21\pi}{3}</math></p> <p>C) <math>\frac{20\pi}{3}</math></p> <p>D) <math>\frac{13\pi}{3}</math></p> <p>E) <math>\frac{31\pi}{3}</math></p>	<p>Reference &amp; Co-terminal Angles.</p>
<p>The reference angle of <math>\theta = 16</math> radians is equal to</p> <p>A) <math>16 - 5\pi</math></p> <p>B) <math>5\pi - 16</math></p> <p>C) <math>16 - 4\pi</math></p> <p>D) <math>4\pi - 16</math></p> <p>E) <math>6\pi - 16</math></p>	<p>Reference Angles.</p>
<p>The reference angle of the angle <math>\theta = 1225^\circ</math> is</p> <p>A) <math>35^\circ</math></p> <p>B) <math>65^\circ</math></p> <p>C) <math>55^\circ</math></p> <p>D) <math>45^\circ</math></p> <p>E) <math>25^\circ</math></p>	<p>Reference Angles.</p>

<p>The reference angle <math>\alpha'</math>, in radians, of the angle <math>\alpha = 920^\circ</math> is equal to:</p> <p>A) <math>\frac{\pi}{9}</math></p> <p>B) <math>\frac{\pi}{3}</math></p> <p>C) <math>\frac{\pi}{5}</math></p> <p>D) <math>\frac{\pi}{10}</math></p> <p>E) <math>\frac{\pi}{6}</math></p>	<p>Reference Angles.</p>
<p>The reference angle of <math>\theta = 2</math> radians is equal to</p> <p>A) <math>\pi - 2</math></p> <p>B) <math>2 - \pi</math></p> <p>C) <math>2 + \pi</math></p> <p>D) <math>2\pi - 2</math></p> <p>E) <math>\frac{\pi}{2} - 2</math></p>	<p>Reference Angles.</p>
<p>The reference angle of <math>-115^\circ</math> is</p> <p>A) <math>65^\circ</math></p> <p>B) <math>55^\circ</math></p> <p>C) <math>75^\circ</math></p> <p>D) <math>45^\circ</math></p> <p>E) <math>25^\circ</math></p>	<p>Reference Angles.</p>

<p>The greatest negative angle that is coterminal with <math>\frac{27\pi}{5}</math> is</p> <p>A) <math>-\frac{3\pi}{5}</math></p> <p>B) <math>-\pi</math></p> <p>C) <math>-\frac{2\pi}{5}</math></p> <p>D) <math>-\frac{4\pi}{5}</math></p> <p>E) <math>-\frac{\pi}{5}</math></p>	<p>Reference &amp; Co-terminal Angles.</p>
<p>If <math>\alpha</math> is the smallest positive coterminal angle of <math>\frac{57\pi}{2}</math> and <math>\beta</math> is reference angle of <math>1270^\circ</math>, then <math>\alpha + \beta =</math></p> <p>A) <math>100^\circ</math></p> <p>B) <math>180^\circ</math></p> <p>C) <math>190^\circ</math></p> <p>D) <math>280^\circ</math></p> <p>E) <math>210^\circ</math></p>	<p>Reference &amp; Co-terminal Angles.</p>
<p>If <math>\alpha</math> is the reference angle of <math>845^\circ</math> and <math>\beta</math> is the least positive coterminal of <math>-705^\circ</math>, then <math>\alpha + \beta =</math></p> <p>A) <math>70^\circ</math></p> <p>B) <math>80^\circ</math></p> <p>C) <math>180^\circ</math></p> <p>D) <math>160^\circ</math></p> <p>E) <math>150^\circ</math></p>	<p>Reference &amp; Co-terminal Angles.</p>

<p>If <math>\theta = \frac{13\pi}{18}</math>, then the degree measure of the reference angle of <math>\theta</math> is</p> <p>(A) <math>50^\circ</math>  B) <math>60^\circ</math>  C) <math>45^\circ</math>  D) <math>70^\circ</math>  E) <math>36^\circ</math></p>	<p>Reference &amp;  Co-terminal  Angles.</p>
<p>If <math>\alpha'</math> is the reference angle of <math>\alpha = -4</math> and <math>\beta'</math> is the reference of <math>\beta = 7</math> then <math>\alpha' + \beta' =</math></p> <p>A) <math>3 - 2\pi</math>  B) <math>3 - \pi</math>  C) <math>11 - 3\pi</math>  D) <math>11 - 2\pi</math>  E) <math>2\pi - 3</math></p>	<p>Reference &amp;  Co-terminal  Angles.</p>
<p>The value of <math>2 - \sin^2(40^\circ) - \sin^2(50^\circ)</math> is</p> <p>A) -3  B) 0  C) 3  D) 1  E) -1</p>	<p>Reference &amp;  Co-terminal  Angles.</p>

<p>The value of <math>1 - \cos^2(20^\circ) - \cos^2(70^\circ)</math> is</p> <p>A) <math>\cos^2(90^\circ)</math>  B) <math>\sin^2(90^\circ)</math>  C) <math>\sin^2(70^\circ)</math>  D) <math>1 - \sin^2(20^\circ)</math>  E) <math>\sin^2(20^\circ) - \cos^2(20^\circ)</math></p>	<p>Co-function Identities.</p>
<p>Let the point <math>(k, -2)</math> lie on the terminal side of angle <math>\theta</math> in standard position. If <math>\csc \theta = -3</math>, where <math>\cos \theta &gt; 0</math>, then the value of <math>k</math> is equal to</p> <p>A) <math>4\sqrt{2}</math>  B) <math>-4\sqrt{2}</math>  C) <math>2\sqrt{2}</math>  D) <math>-2\sqrt{2}</math>  E) <math>-\sqrt{2}</math></p>	<p>Trigonometric Functions of Angles.</p>
<p>If the terminal side of an angle <math>\theta</math>, in standard position, is in quadrant III and has slope equal <math>\frac{1}{2}</math>, then <math>\sin \theta + \cos \theta =</math></p> <p>A) <math>-\frac{3\sqrt{5}}{5}</math>  B) <math>-\frac{\sqrt{5}}{5}</math>  C) <math>\sqrt{5}</math>  D) <math>-\frac{2\sqrt{3}}{5}</math>  E) <math>\sqrt{3}</math></p>	<p>Trigonometric Functions of Angles</p>

<p>If <math>\tan \theta = 4</math> and <math>P(-3, n)</math> is a point on the terminal side of <math>\theta</math> where <math>\theta</math> is in standard position, then <math>\sec \theta =</math></p> <p>A) <math>\sqrt{17}</math>  B) <math>-\frac{5}{3}</math>  C) <math>-\sqrt{17}</math>  D) <math>-\frac{1}{4}</math>  E) <math>-\frac{\sqrt{17}}{4}</math></p>	<p>Trigonometric Functions of Angles</p>
<p>If the terminal side of an angle <math>\theta</math> in standard position is given by <math>Ax + y = 0</math>, <math>x &lt; 0</math> and <math>\sin \theta = \frac{1}{3}</math>, then <math>A =</math></p> <p>A) <math>\frac{\sqrt{2}}{4}</math>  B) <math>-\frac{\sqrt{2}}{4}</math>  C) <math>\frac{3\sqrt{2}}{2}</math>  D) 1  E) <math>\frac{3\sqrt{2}}{8}</math></p>	<p>Trigonometric Functions of Angles</p>

<p>If the point <math>\left(-\frac{1}{2}, \frac{\sqrt{3}}{2}\right)</math> is on the terminal side of the angle <math>\theta</math> in standard position, then <math>\tan \theta =</math></p> <p>A) <math>-\frac{\sqrt{3}}{3}</math></p> <p>B) <math>-2</math></p> <p><b>C) <math>-\sqrt{3}</math></b></p> <p>D) <math>-\frac{2\sqrt{3}}{3}</math></p> <p>E) <math>-\frac{\sqrt{3}}{2}</math></p>	<p>Trigonometric Functions of Angles</p>
<p>If the terminal side of an angle <math>\theta</math>, in standard position, is defined by <math>x - 2y = 0, x &gt; 0</math>, then <math>\sec \theta =</math></p> <p>A) <math>\frac{2\sqrt{5}}{5}</math></p> <p><b>B) <math>\frac{\sqrt{5}}{2}</math></b></p> <p>C) <math>\frac{1}{2}</math></p> <p>D) <math>\sqrt{5}</math></p> <p>E) <math>\frac{5}{2}</math></p>	<p>Trigonometric Functions of Angles</p>
<p>If the terminal side of an angle <math>\theta</math> in standard position is defined by <math>3x + 2y = 0, x \leq 0</math>, then <math>\csc \theta =</math></p> <p><b>A) <math>\frac{\sqrt{13}}{3}</math></b></p> <p>B) <math>-\sqrt{13}</math></p> <p>C) <math>\frac{3\sqrt{13}}{13}</math></p> <p>D) <math>-\frac{3\sqrt{13}}{13}</math></p> <p>E) <math>-\frac{\sqrt{13}}{3}</math></p>	<p>Trigonometric Functions of Angles</p>

<p>If the terminal side of an angle <math>\theta</math> in standard position is given by <math>3x - y = 0, x &lt; 0</math>, then <math>\csc \theta =</math></p> <p>A) <math>-\frac{3\sqrt{10}}{10}</math></p> <p>B) <math>\frac{\sqrt{10}}{3}</math></p> <p>C) <math>\frac{3\sqrt{10}}{10}</math></p> <p>D) <math>-\frac{\sqrt{10}}{3}</math></p> <p>E) -3</p>	<p>Trigonometric Functions of Angles</p>
<p>Suppose that the terminal side of the angle <math>\theta</math> in standard position is given by <math>12x - 5y = 0, x \leq 0</math>, then <math>\frac{60}{13}(\sec \theta + \csc \theta) =</math></p> <p>A) -17</p> <p>B) -7</p> <p>C) 7</p> <p>D) -8</p> <p>E) 17</p>	<p>Trigonometric Functions of Angles</p>
<p>If the equation of the terminal side of <math>\theta</math> in standard position is <math>x + 2y = 0, x \geq 0</math>, then <math>\sin \theta - \cos \theta =</math></p> <p>A) <math>-\frac{3\sqrt{5}}{5}</math></p> <p>B) <math>-\frac{\sqrt{5}}{5}</math></p> <p>C) <math>-\sqrt{5}</math></p> <p>D) <math>\frac{\sqrt{5}}{5}</math></p> <p>E) <math>\sqrt{5}</math></p>	<p>Trigonometric Functions of Angles</p>



<p>If the equation of the terminal side of an angle <math>\theta</math> in standard position is <math>4x + 3y = 0</math>, where <math>x &lt; 0</math>, then <math>\csc \theta + \sec \theta =</math></p> <p>A) <math>-\frac{5}{12}</math></p> <p>B) <math>\frac{5}{12}</math></p> <p>C) <math>-\frac{7}{12}</math></p> <p>D) <math>\frac{7}{5}</math></p> <p>E) <math>-\frac{1}{5}</math></p>	<p>Trigonometric Functions of Angles</p>
<p>If the point <math>(2, -3)</math> is on the terminal side of the angle <math>\theta</math> in the standard position, then <math>12\csc \theta + 4\sec \theta</math> is equal to</p> <p>A) <math>-2\sqrt{13}</math></p> <p>B) <math>-\sqrt{13}</math></p> <p>C) <math>\sqrt{13}</math></p> <p>D) <math>2\sqrt{13}</math></p> <p>E) <math>-\frac{18\sqrt{13}}{13}</math></p>	<p>Trigonometric Functions of Angles</p>
<p>The equation of the terminal side of <math>\theta</math> is given by <math>\sqrt{3}x + y = 0</math>, where <math>x \leq 0</math>, then <math>\csc \theta =</math></p> <p>A) <math>\frac{2\sqrt{3}}{3}</math></p> <p>B) <math>\frac{\sqrt{3}}{2}</math></p> <p>C) <math>-\frac{\sqrt{3}}{3}</math></p> <p>D) <math>-\frac{2\sqrt{3}}{3}</math></p> <p>E) <math>\frac{1}{2}</math></p>	<p>Trigonometric Functions of Angles</p>

<p>If the equation of the terminal side of an angle <math>\theta</math> in standard position is <math>4x - 3y = 0, x &lt; 0</math>, then <math>4\csc \theta + 9\tan \theta =</math></p> <p>A) 7  B) -5  C) 11  D) -6  E) 10</p>	<p>Trigonometric Functions of Angles</p>
<p>If the terminal side of the angle <math>\theta</math> in the standard position coincides with the line <math>\sqrt{3}x + y = 0</math>, with <math>x \leq 0</math> then</p> <p>(a) <math>\cot \theta = -\frac{\sqrt{3}}{3}</math>  (b) <math>\tan \theta = \sqrt{3}</math>  (c) <math>\sin \theta = -\frac{1}{2}</math>  (d) <math>\cos \theta = \frac{\sqrt{3}}{2}</math>  (e) <math>\tan \theta = -2</math></p>	<p>Trigonometric Functions of Angles</p>
<p>If <math>12x - 5y = 0, x \leq 0</math>, is the equation of the terminal side of an angle <math>\alpha</math>, then <math>5\tan \alpha - 12\csc \alpha =</math></p> <p>A) 25  B) - 1  C) 15  D) 20  E) -25</p>	<p>Trigonometric Functions of Angles</p>

<p>Suppose that the terminal side of an angle <math>\theta</math> in standard position lies on the line <math>y = -\frac{1}{2}x</math> where <math>x &gt; 0</math>, then <math>\sin \theta + \tan \theta =</math></p> <p>A) <math>-\frac{5+2\sqrt{5}}{10}</math></p> <p>B) <math>-\frac{10+\sqrt{5}}{5}</math></p> <p>C) <math>\frac{2-\sqrt{5}}{5}</math></p> <p>D) <math>-\frac{10-2\sqrt{5}}{10}</math></p> <p>E) <math>\frac{1-\sqrt{5}}{10}</math></p>	<p>Trigonometric Functions of Angles</p>
<p>If <math>\sin \theta = \frac{1}{5}</math>, and <math>p(-3, k)</math> is a point on the terminal side of <math>\theta</math> in standard position, then the value of <math>k</math> is:</p> <p>A) <math>\frac{\sqrt{6}}{4}</math></p> <p>B) <math>-\frac{\sqrt{6}}{2}</math></p> <p>C) <math>-\frac{\sqrt{6}}{4}</math></p> <p>D) 1</p> <p>E) <math>\frac{\sqrt{6}}{2}</math></p>	<p>Trigonometric Functions of Angles</p>
<p>If the terminal side of the angle <math>\theta</math> in standard position is defined by <math>6x + 8y = 0, y &lt; 0</math> then <math>10\cos \theta - 12\tan \theta =</math></p> <p>A) 17</p> <p>B) -17</p> <p>C) -1</p> <p>D) 1</p> <p>E) 24</p>	<p>Trigonometric Functions of Angles</p>

If the angle  $\theta = 12$  radian, then

(a)  $\theta$  is in the fourth quadrant

(b)  $\theta$  is a quadrantal angle

(c)  $\theta$  is in the first quadrant

(d)  $\theta$  is in the second quadrant

(e)  $\theta$  is in the third quadrant

Angles in  
radians