

### 6.3: (Trigonometric Graphs)

<p>If <math>y = -2 - 3\sin\left(2x + \frac{2\pi}{3}\right)</math>, M is the maximum, P the period and S the phase shift then <math>M + \frac{P+S}{\pi}</math> is equal to</p> <p>A) <math>\frac{7}{3}</math>          B) <math>\frac{-1}{3}</math>          C) <math>\frac{4}{3}</math>          D) <math>\frac{17}{3}</math>          E) <math>\frac{5}{3}</math></p>	<p>Trigonometric Graphs (<math>\sin \theta</math>).</p>
<p>The graph of <math>y = -\sin(\pi x + \pi)</math>, <math>-1 \leq x \leq 1</math>, is increasing on the interval:</p> <p>A) <math>\left(-\frac{1}{2}, \frac{1}{2}\right)</math>          B) <math>\left(-1, \frac{1}{2}\right)</math>          C) <math>\left(-\frac{1}{2}, 1\right)</math>          D) <math>(0, 1)</math>          E) <math>\left(\frac{1}{2}, 1\right)</math></p>	<p>Trigonometric Graphs (<math>\sin \theta</math>).</p>
<p>The graph of <math>y = -\frac{2}{3}\sin\frac{\pi}{4}x</math>, with <math>-8 \leq x \leq 0</math>,</p> <p>A) is above the <math>x</math>-axis on the interval <math>(-4, 0)</math>          B) intersects the <math>x</math>-axis at two points          C) has maximum value of <math>2/3</math> in the interval <math>[-8, -4]</math>          D) is increasing on the intervals <math>(-8, -6)</math> and <math>(-2, 0)</math>          E) is decreasing on the interval <math>(-6, -2)</math></p>	<p>Trigonometric Graphs (<math>\sin \theta</math>).</p>

<p>The graph of the function <math>f(x) = -3\sin\left(\frac{1}{3}x\right)</math>, with <math>0 \leq x \leq 6\pi</math>, lies completely below the <math>x</math>-axis on the interval</p> <p>(a) <math>(0, 3\pi)</math></p> <p>(b) <math>(0, 6\pi)</math></p> <p>(c) <math>(3\pi, 6\pi)</math></p> <p>(d) <math>(4\pi, 6\pi)</math></p> <p>(e) <math>(\pi, 6\pi)</math></p>	<p>Trigonometric Graphs (<math>\sin \theta</math>).</p>
<p>Which one of the following statements is FALSE?</p> <p>A) <math>f(x) = x + \cos x \sin x</math> is an odd function.</p> <p>B) If <math>\frac{\pi}{2} &lt; x &lt; \pi</math>, then <math>\sin x - \cos x &gt; 0</math>.</p> <p>C) <math>f(x) = \cos x \sin^2 x</math> is an odd function.</p> <p>D) <math>f(x) = 3 + \cos x</math> is an even function.</p> <p>E) The period of the function <math>f(x) = -\sin(2\pi x)</math> is 1</p>	<p>Trigonometric Graphs (<math>\sin \theta</math>).</p>
<p>The graph of the function <math>y = 3\sin  x </math>, <math>-\pi \leq x \leq \frac{\pi}{2}</math>, is decreasing on the interval</p> <p>A) <math>(-\pi, 0)</math></p> <p>B) <math>\left(-\frac{\pi}{2}, \frac{\pi}{2}\right)</math></p> <p>C) <math>\left(-\frac{\pi}{2}, 0\right)</math></p> <p>D) <math>\left(-\pi, -\frac{\pi}{2}\right)</math></p> <p>E) <math>\left(0, \frac{\pi}{2}\right)</math></p>	<p>Trigonometric Graphs (<math>\sin \theta</math>).</p>

<p>The graph of the function <math>f(x) = -3\sin\left(\frac{\pi}{2} - 2x\right)</math>, <math>\frac{\pi}{4} \leq x \leq \frac{5\pi}{4}</math> is below the <math>x</math> - axis on</p> <p>A) <math>\left(\frac{3\pi}{4}, \frac{5\pi}{4}\right)</math></p> <p>B) <math>\left(\frac{\pi}{2}, \pi\right)</math></p> <p>C) <math>\left(\frac{\pi}{2}, \frac{3\pi}{4}\right)</math></p> <p>D) <math>\left(\frac{\pi}{4}, \pi\right)</math></p> <p>E) <math>\left(\frac{\pi}{4}, \frac{\pi}{2}\right)</math></p>	<p>Trigonometric Graphs (<math>\sin \theta</math>).</p>
<p>The range of the function <math>f(x) = 1 + \sqrt{4 - 4\sin^2 x}</math>, is</p> <p>A) [1,5]</p> <p>B) [0,2]</p> <p>C) [-1,3]</p> <p>D) [1,3]</p> <p>E) [-3,5]</p>	<p>Trigonometric Graphs (<math>\sin \theta</math>).</p>
<p>If the range of the function <math>y = K - 5\sin\left(\frac{\pi}{2}x - \frac{\pi}{2}\right)</math> is <math>[-7,3]</math>, then <math>K =</math></p> <p>A) -2</p> <p>B) <math>\pi + 2</math></p> <p>C) 0</p> <p>D) 4</p> <p>E) <math>2 - \pi</math></p>	<p>Trigonometric Graphs (<math>\sin \theta</math>).</p>

<p>The graph of <math>y = 3\sin  \pi x </math>, with <math>-1 \leq x \leq 1</math>, is increasing on the interval</p> <p>A) <math>\left(-1, -\frac{1}{2}\right) \cup \left(0, \frac{1}{2}\right)</math></p> <p>B) <math>\left(-\frac{1}{2}, 0\right) \cup \left(\frac{1}{2}, 1\right)</math></p> <p>C) <math>\left(-1, \frac{1}{2}\right)</math></p> <p>D) <math>\left(-\frac{1}{2}, \frac{1}{2}\right)</math></p> <p>E) <math>\left(-\frac{1}{2}, 1\right)</math></p>	<p>Trigonometric Graphs (<math>\sin \theta</math>).</p>
<p>Which one of the following statements is FALSE about the graph of the function <math>f(x) = -\frac{1}{2}\sin\left(\frac{\pi x}{2}\right)</math>, <math>-2 \leq x \leq 2</math> ?</p> <p>A) is increasing on the interval <math>(-1, 1)</math>.</p> <p>B) has three <math>x</math>-intercepts.</p> <p>C) lies above the <math>x</math>-axis on the interval <math>(-2, 0)</math>.</p> <p>D) lies below the <math>x</math>-axis on the interval <math>(0, 2)</math>.</p> <p>E) passes through the origin.</p>	<p>Trigonometric Graphs (<math>\sin \theta</math>).</p>
<p>The graph of the function <math>f(x) = 2\sin(4x + \pi)</math>, where <math>-\frac{\pi}{4} \leq x \leq \frac{\pi}{4}</math>,</p> <p>A) has three <math>x</math>-intercepts</p> <p>B) has no <math>y</math>-intercept</p> <p>C) is increasing on the interval <math>\left(-\frac{\pi}{8}, \frac{\pi}{8}\right)</math></p> <p>D) is below the <math>x</math>-axis in the interval <math>\left(-\frac{\pi}{4}, 0\right)</math></p> <p>E) is above the <math>x</math>-axis in the interval <math>\left(0, \frac{\pi}{4}\right)</math></p>	<p>Trigonometric Graphs (<math>\sin \theta</math>).</p>

<p>Which one of the following statements is TRUE about the function <math>f(x) = -1 + \frac{1}{2} \sin(2x - 3\pi)</math>?</p> <p>A) The range of the function is <math>\left[-\frac{3}{2}, -\frac{1}{2}\right]</math></p> <p>B) The vertical translation of the graph of the function is <math>\frac{1}{2}</math> unit up</p> <p>C) The period of the function is <math>2\pi</math></p> <p>D) The phase shift of the function is <math>3\pi</math> unit to the right</p> <p>E) The amplitude of the function is <math>\frac{3}{2}</math></p>	<p>Trigonometric Graphs (<math>\sin \theta</math>).</p>
<p>The graph of <math>y = \frac{5}{2} - \cos\left[3\left(x - \frac{\pi}{6}\right)\right]</math> has:</p> <p>A) range <math>\left[\frac{3}{2}, \frac{7}{2}\right]</math></p> <p>B) period <math>\pi</math></p> <p>C) phase shift <math>\frac{\pi}{2}</math> to the left</p> <p>D) amplitude 3</p> <p>E) <math>y</math>-intercept <math>-\frac{5}{2}</math></p>	<p>Trigonometric Graphs (<math>\cos \theta</math>).</p>
<p>The number of <math>x</math>-intercepts of the graph of <math>y = -2\cos \pi x, -\frac{3}{2} \leq x &lt; \frac{5}{2}</math>. is</p> <p>A) 5</p> <p>B) 1</p> <p>C) 3</p> <p>D) 4</p> <p>E) 2</p>	<p>Trigonometric Graphs (<math>\cos \theta</math>).</p>


The range of the function  $y = 2 - |3\cos(3x - \pi)|$ , is

- A)  $[-1, 2]$
- B)  $[-3, 0]$
- C)  $(-\infty, -3] \cup [0, \infty)$
- D)  $[1, 4]$
- E)  $(-\infty, 1] \cup [4, \infty)$

Trigonometric  
Graphs  
( $\cos \theta$ ).

If the adjacent figure represents the graph of  $y = -2\cos(bx + c)$ , then

- A)  $b = \pi$  and  $c = \frac{\pi}{2}$
- B)  $b = \pi$  and  $c = -\frac{\pi}{2}$
- C)  $b = 2\pi$  and  $c = -\frac{\pi}{4}$
- D)  $b = 2\pi$  and  $c = \frac{\pi}{4}$
- E)  $b = \pi$  and  $c = \frac{\pi}{4}$



Trigonometric  
Graphs  
( $\cos \theta$ ).

The graph of the function  $f(x) = 3\cos(2\pi x)$  over  $[-1, 1]$  intersects the line  $y = -2$  at

- A) 5 points
- B) 7 points
- C) 8 points
- D) 3 points
- E) 4 points

Trigonometric  
Graphs  
( $\cos \theta$ ).

The graph of the function  $f(x) = -2\cos\left(\frac{\pi}{4}x - \frac{\pi}{2}\right)$ ,  $0 \leq x \leq 10$  is increasing on the interval

- A) [2,6]
- B) [0,4]
- C)  $[0,4] \cup [8,10]$
- D) [8,10]
- E)  $[3,4] \cup [9,10]$

Trigonometric Graphs ( $\cos \theta$ ).

If the adjacent figure represents the graph of the function  $y = a\cos(kx + b)$ , then  $a + k + \frac{b}{\pi} =$


- A) 0
- B) 1
- C) -1
- D)  $\frac{\pi-1}{2}$
- E)  $\frac{\pi^2-1}{2}$



Trigonometric Graphs ( $\cos \theta$ ).

If the adjacent figure represents the graph of  $y = a\cos(bx) + 1$ , then  $2b - \pi a =$

- A)  $3\pi$
- B)  $2\pi$
- C)  $-2\pi$
- D)  $-3\pi$
- E)  $\pi$




Trigonometric Graphs ( $\cos \theta$ ).

<p>If the graph of the function <math>f(x) = 2\cos(bx + c)</math>, <math>b &gt; 0</math> and <math>-\frac{\pi}{2} \leq c \leq \frac{\pi}{2}</math>, has period <math>\frac{2\pi}{3}</math> and passes through the point <math>(0,2)</math>, then <math>b + c =</math></p> <p><b>A) 3</b>  B) 0  C) 4  D) 5  E) 2</p>	<p>Trigonometric Graphs (<math>\cos \theta</math>).</p>
<p>The range of the function <math>f(x) = 3 -  \cos(x - \pi) </math> is</p> <p><b>A) [2,3]</b>  B) [2,4]  C) [-3,3]  D) (2,4)  E) (2,3)</p>	<p>Trigonometric Graphs (<math>\cos \theta</math>).</p>
<p>Which one of the following statements is FALSE about the function <math>f(x) = 5\cos\left(3x - \frac{\pi}{4}\right)</math> ?</p> <p><b>(a) The graph of <math>f(x)</math> has <math>y</math>-intercept at <math>y = -1</math></b>  (b) The range of <math>f(x)</math> is <math>[-5,5]</math>  (c) The horizontal shift of the graph of <math>f(x)</math> is <math>\frac{\pi}{12}</math> units to the right  (d) The amplitude of the graph of <math>f(x)</math> is 5  (c) The period of <math>f(x)</math> is <math>\frac{2\pi}{3}</math></p>	<p>Trigonometric Graphs (<math>\cos \theta</math>).</p>



<p>The graph of the function <math>f(x) = -2\sin(3x)</math> over <math>[0, 2\pi]</math> intersects the line <math>y = \frac{3}{2}</math> at</p> <p>A) 3 points.            B) 7 points.  <b>C) 6 points.</b>            D) 2 points.            E) 5 points</p>	<p>Trigonometric Graphs (<math>\sin \theta</math>).</p>
<p>The equation of the function whose part of its graph is drawn below is</p> <p>A) <math>y = -\cos 4x</math>            B) <math>y = -\sin 3x</math>            C) <math>y = -\sin 2x</math>  <b>D) <math>y = \cos 2x</math></b>            E) <math>y = -\cos 2x</math></p>	<p>Trigonometric Graphs (<math>\cos \theta</math>).</p>
<p>If A is the amplitude, P is the period and F is the phase shift of the graph of the function <math>f(x) = -3\cos\left(\frac{\pi x}{5} - 2\pi\right)</math>, then <math>\frac{-2A+P-F}{2} =</math></p> <p>A) <math>-3 - 10\pi</math>  <b>B) -3</b>            C) 3            D) <math>3 + 10\pi</math>            E) <math>-3 + 10\pi</math></p>	<p>Trigonometric Graphs (<math>\cos \theta</math>).</p>
<p>The range <math>R</math> and the period <math>P</math> of the function <math>y = -\left 3\sin\frac{x}{2}\right </math> are given by</p> <p><b>(a) <math>R = [-3, 0]</math>, <math>P = 2\pi</math></b>            (b) <math>R = [-3, 3]</math>, <math>P = 2\pi</math>            (c) <math>R = [-3, 0]</math>, <math>P = 4\pi</math>            (d) <math>R = [-3, 0]</math>, <math>P = \pi</math>            (e) <math>R = [-3, 0]</math>, <math>P = \frac{\pi}{2}</math></p>	<p>Trigonometric Graphs (<math>\sin \theta</math>).</p>



<p>The graph of <math>y = -\frac{1}{2}\sin \pi x</math></p> <p>A) has a maximum of <math>\frac{1}{2}</math> in the interval <math>[3,4]</math></p> <p>B) is decreasing on the interval <math>[1,2]</math></p> <p>C) is increasing on the interval <math>[0,1]</math></p> <p>D) has a minimum of <math>-\frac{1}{2}</math> in the interval <math>[-1,0]</math></p> <p>E) has a minimum of <math>-\frac{1}{2}</math> in the interval <math>[1,2]</math></p>	<p>Trigonometric Graphs (<math>\sin \theta</math>).</p>
<p>The number of points of intersection of <math>y = 1</math> and <math>y = -2\sin \frac{\pi x}{3}</math> in the interval <math>[-3,9]</math></p> <p>A) 2</p> <p>B) 4</p> <p>C) 5</p> <p>D) 3</p> <p>E) 6</p>	<p>Trigonometric Graphs (<math>\sin \theta</math>).</p>
<p>Let <math>f(x) = -\frac{1}{3}\cos(2\pi x)</math> and <math>A</math> be its amplitude, <math>P</math> be its period and <math>(a, b)</math> be the highest point of the function in the interval <math>[0,1]</math>. Then, <math>A + P + a - b =</math></p> <p>A) <math>\frac{3}{2}</math></p> <p>B) <math>\frac{5}{2}</math></p> <p>C) <math>\frac{3\pi}{2}</math></p> <p>D) <math>1 + \frac{3\pi}{2}</math></p> <p>E) <math>\frac{5\pi-1}{2}</math></p>	<p>Trigonometric Graphs (<math>\cos \theta</math>).</p>

The range of the function  $f(x) = -\left|3\sin\frac{2x}{3}\right|$  is

A)  $[-3,0]$

B)  $[0,3]$

C)  $[-3,3]$

D)  $[-1,0]$

E)  $[-1,1]$

Trigonometric  
Graphs  
( $\sin \theta$ ).

The function  $y = \cos\left(\frac{x}{3} - \frac{\pi}{3}\right)$ , for  $-\frac{13\pi}{2} \leq x \leq \frac{17\pi}{2}$  has

A) three maximum values

B) two maximum values

C) four maximum values

D) five maximum values

E) six maximum values

Trigonometric  
Graphs  
( $\sin \theta$ ).

If the adjacent figure represents the graph of the function  $y = a\sin(bx + c)$ ,

$-\frac{3\pi}{2} \leq x \leq \frac{5\pi}{2}$ , then


A)  $a = -3, b = \frac{1}{2}, c = \frac{3\pi}{4}$

B)  $a = -3, b = 2, c = -\frac{3\pi}{2}$

C)  $a = 3, b = \frac{1}{2}, c = -\frac{3\pi}{4}$

D)  $a = 3, b = 4, c = 3\pi$

E)  $a = -3, b = \frac{1}{4}, c = \frac{3\pi}{8}$



Trigonometric  
Graphs  
( $\sin \theta$ ).


The graph of the function  $y = \frac{1}{2} \cos \frac{\pi}{2} x$  increases over the interval

- A) (2,4)
- B) (0,2)
- C) (3,5)
- D) (1,3)
- E) (2,5)

Trigonometric  
Graphs  
( $\cos \theta$ ).

The adjacent figure represents part of the graph of

- A)  $y = \sin\left(\frac{1}{2}x\right) + 1$
- B)  $y = \cos\left(\frac{1}{2}x\right) + 1$
- C)  $y = 2\sin(x) + 1$
- D)  $y = 2\cos(x) + 1$
- E)  $y = \sin(2x) + 1$



Trigonometric  
Graphs  
( $\sin \theta$ ).

The graph of  $y = -|\sin \pi x|$  over the interval  $\left[\frac{1}{2}, \frac{3}{2}\right]$  intersects the line  $y = -\frac{1}{2}$  at:

- (a) 2 points
- (b) 1 point
- (c) 3 points
- (d) 4 points
- (e) no point

Trigonometric  
Graphs  
( $\sin \theta$ ).

If the adjacent figure represents the graph of the function  $y = a\cos(bx + c)$ ,  $-\frac{2\pi}{3} \leq x \leq \frac{2\pi}{3}$  then


A)  $a = -2, b = 3, c = \pi$

B)  $a = 2, b = \frac{1}{3}, c = \pi$

C)  $a = -2, b = 3, c = -\frac{2\pi}{3}$

D)  $a = -2, b = \frac{2}{3}, c = \frac{2\pi}{3}$

E)  $a = 2, b = \frac{2}{3}, c = \frac{\pi}{3}$



Trigonometric  
Graphs  
( $\cos \theta$ ).

Over the interval  $\left[-\frac{\pi}{4}, \frac{15\pi}{4}\right)$ , the function  $y = -\sin\left(\frac{x}{2} + \frac{\pi}{8}\right)$  is decreasing on the interval(s)

A)  $\left[-\frac{\pi}{4}, \frac{3\pi}{4}\right]$  and  $\left[\frac{11\pi}{4}, \frac{15\pi}{4}\right]$

B)  $\left[\frac{3\pi}{4}, \frac{11\pi}{4}\right]$

C)  $\left[-\frac{\pi}{4}, \frac{5\pi}{4}\right]$  and  $\left[\frac{11\pi}{4}, \frac{15\pi}{4}\right]$

D)  $\left[0, \frac{3\pi}{4}\right]$  and  $\left[\frac{5\pi}{4}, \frac{11\pi}{4}\right]$

E)  $\left[-\frac{\pi}{4}, \frac{3\pi}{4}\right]$  and  $\left[\frac{7\pi}{4}, \frac{11\pi}{4}\right]$

Trigonometric  
Graphs  
( $\sin \theta$ ).


The function  $y = -2 + 5\cos(2x + 3\pi)$  has

- A) range =  $[-7,3]$ , phase shift =  $\frac{3\pi}{2}$  units to the left
- B) range =  $[-5,5]$  phase shift =  $3\pi$  units to the left
- C) range =  $[-7,3]$  phase shift =  $\frac{3\pi}{2}$  units to the right
- D) range =  $[-2,2]$ , phase shift =  $3\pi$  units to the right
- E) range =  $[-7,7]$ , phase shift =  $\frac{3\pi}{2}$  units to the left

Trigonometric  
Graphs  
( $\cos \theta$ ).

$y = -2\cos(bx + c)$ , then

- A)  $b = \frac{\pi}{2}$  and  $c = \frac{\pi}{2}$
- B)  $b = \frac{\pi}{2}$  and  $c = \frac{\pi}{4}$
- C)  $b = \pi$  and  $c = +\frac{\pi}{2}$
- D)  $b = 2\pi$  and  $c = -\frac{\pi}{4}$
- E)  $b = 2\pi$  and  $c = -\pi$



Trigonometric  
Graphs  
( $\cos \theta$ ).


Which one of the following statements is TRUE?

- A) The equation  $\sin x = 2$  has no solution in the interval  $\frac{\pi}{2} \leq x \leq \frac{3\pi}{2}$ .
- B) Domain of  $y = \sin(x + \pi)$  is  $[-\pi, \infty)$ .
- C) The period of the graph of  $y = \sin[\pi(2x - 1)]$  is  $\pi$ .
- D) The range of the function  $y = 2\sin(x - \pi) - 1$  is  $[-2,2]$ .
- E)  $(\sec x)(\sin x) = 1$ .

Trigonometric  
Graphs  
( $\sin \theta$ ).

$y = \frac{1}{2} \cos(bx + c), \frac{\pi}{2} \leq x \leq \frac{9\pi}{2}$  then

- A)  $b = \frac{1}{2}, c = -\frac{\pi}{4}$
- B)  $b = \frac{1}{2}, c = -\frac{\pi}{2}$
- C)  $b = 2, c = -\frac{\pi}{2}$
- D)  $b = \frac{1}{4}, c = \frac{\pi}{2}$
- E)  $b = \frac{1}{4}, c = \frac{\pi}{4}$



Trigonometric  
Graphs  
( $\cos \theta$ ).

The graph of the function  $y = 2\sin\frac{1}{4}x$ , for  $-8\pi \leq x \leq 8\pi$ , has

- A) two maximum values and five  $x$ -intercepts
- B) three maximum values and four  $x$ -intercepts
- C) four maximum values and three  $x$ -intercepts
- D) five maximum values and two  $x$ -intercepts
- E) three maximum values and three  $x$ -intercepts

Trigonometric  
Graphs  
( $\sin \theta$ ).

For  $-8 \leq x \leq 8$ , the graph of the function  $y = -\frac{3}{2} \sin\left(\frac{\pi}{4}x\right)$  lies below the  $x$ -axis in the interval(s) [Hint: Sketch the graph]

- A)  $(-8, -4)$  and  $(0, 4)$
- B)  $(-4, 0)$  and  $(4, 8)$
- C)  $(-6, -2)$  and  $(2, 6)$
- D)  $(-4, 0)$  and  $(0, 4)$
- E)  $(-4, 4)$

Trigonometric  
Graphs  
( $\sin \theta$ ).


The graph of the function  $y = 3\cos\left(\frac{1}{2}x - \frac{\pi}{4}\right)$ , on the interval  $\left[-\frac{7\pi}{2}, \frac{9\pi}{2}\right]$  intersects the line  $y = 2$  at

- A) four points
- B) six points
- C) three points
- D) five points
- E) no points

Trigonometric Graphs ( $\cos \theta$ ).

$y = -2\cos(bx + c)$ ,  $-\frac{2\pi}{3} \leq x \leq \frac{2\pi}{3}$  and  $b > 0$ , then


- A)  $b = 3, c = \pi$
- B)  $b = \frac{1}{3}, c = \pi$
- C)  $b = 3, c = -\frac{2\pi}{3}$
- D)  $b = \frac{2}{3}, c = \frac{2\pi}{3}$
- E)  $b = \frac{2}{3}, c = \frac{\pi}{3}$



Trigonometric Graphs ( $\cos \theta$ ).

If the adjacent figure represents the graph of  $y = a\cos(bx + c)$  over one period where  $0 < c \leq 2\pi$  and  $a < 0$ , then  $8a + b + c =$

- A)  $2\pi$
- B)  $-2\pi$
- C)  $\frac{\pi}{2}$
- D)  $-\pi$
- E)  $\pi$



Trigonometric Graphs ( $\cos \theta$ ).




The graph of  $y = -4\sin(2x - \pi)$  is below  $x$ -axis on the interval :

- A)  $(\frac{\pi}{2}, \pi)$
- B)  $(0, \pi)$
- C)  $(-\pi, -\frac{\pi}{2})$
- D)  $(0, \frac{\pi}{2})$
- E)  $(-\frac{\pi}{2}, \frac{\pi}{2})$

Trigonometric  
Graphs  
( $\sin \theta$ ).

The adjacent graph represents a part of the graph of

- A)  $y = \sin(x + \frac{\pi}{4})$
- B)  $y = \sin(x - \frac{\pi}{4})$
- C)  $y = \cos(x + \frac{\pi}{4})$
- D)  $y = 1 - \cos(x - \frac{\pi}{4})$
- E)  $y = -1 - \sin(x - \frac{\pi}{4})$



Trigonometric  
Graphs  
( $\sin \theta$ ).

If  $f(x) = a\sin bx$ ,  $b > 0$  has a period of  $\frac{2\pi}{3}$  and  $f(\frac{\pi}{2}) = 2$ , then  $a$  is:

- A) -2
- B) 2
- C)  $2\pi$
- D)  $-\pi$
- E)  $\pi$

Trigonometric  
Graphs  
( $\sin \theta$ ).

The graph of  $y = \frac{5}{2} - \cos\left[3\left(x - \frac{\pi}{6}\right)\right]$  has:

A) range  $\left[\frac{3}{2}, \frac{7}{2}\right]$

B) period  $\pi$

C) phase shift  $\frac{\pi}{2}$  to the left.

D) amplitude 3

E) y-intercept  $-\frac{5}{2}$

Trigonometric  
Graphs  
( $\cos \theta$ ).